

PRODUCT INFORMATION

Common Name	H8, mAb5T4
Synonyms	M6P1;WAIF1;TPBG
Applications	ELISA; Flow Cyt
Recommended Dilutions	ELISA 1:5000-10000; Flow Cyt 1:100
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Host Species	Humanized
IgG type	IgG1
Reactivity	Human
Target	5T4
Uniprot ID	Q13641
Description	Anti-5T4 (H8 biosimilar) mAb
Delivery	In Stock
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
Usage	Research use only
Conjugate	Unconjugated






Figure 1. 5T4 protein is highly expressed on the surface of Expi293 cell membrane. Flow cytometry analysis with 15 µg/mL Anti-5T4 (H8 biosimilar) mAb (BME100157) on Expi293 cells transfected with human 5T4 (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram), and Isotype antibody on Expi293 transfected with irrelevant protein (Orange histogram)

Anti-5T4 (H8 biosimilar) mAb ELISA

0.2 µg of Human 5T4, His tagged protein per well




Figure 2. ELISA plate pre-coated by 2 µg/mL (100 µL/well) Human 5T4 Protein, His Tag (PME100123) can bind Anti-5T4 (H8 biosimilar) mAb (BME100157) in a linear range of 0.64–16 ng/mL.

