

PRODUCT INFORMATION

Clone ID	2F4
Target	AGR2
Synonyms	AG-2;hAG-2;HPC8
Host Species	Rabbit
Description	Anti-AGR2 antibody(2F4), Rabbit mAb
Delivery	In Stock
Uniprot ID	O95994
IgG type	Rabbit mAb
Clonality	Monoclonal
Reactivity	Human
Applications	ELISA
Recommended Dilutions	ELISA 1/5000-10000
Purification	Purified from cell culture supernatant by affinity chromatography
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene encodes a member of the disulfide isomerase (PDI) family of endoplasmic reticulum (ER) proteins that catalyze protein folding and thiol-disulfide interchange reactions. The encoded protein has an N-terminal ER-signal sequence, a catalytically active thioredoxin domain, and a C-terminal ER-retention sequence. This protein plays a role in cell migration, cellular transformation and metastasis and is as a p53 inhibitor. As an ER-localized molecular chaperone, it plays a role in the folding, trafficking, and assembly of cysteine-rich transmembrane receptors and the cysteine-rich intestinal glycoprotein mucin. This gene has been implicated in inflammatory bowel disease and cancer progression. [provided by RefSeq, Mar 2017]
Usage	Research use only
Conjugate	Unconjugated






Figure 1. ELISA plate pre-coated by 1 µg/ml (100 µl/well) Human AGR2 Protein can bind Rabbit anti-AGR2 monoclonal antibody(clone: 2F4) in a linear range of 1-50 ng/ml.

DIMABIO CONFIDENTIAL

