


**PRODUCT INFORMATION**

<b>Clone ID</b>	DM205
<b>Target</b>	ANGPTL3
<b>Synonyms</b>	ANG-5; ANGPT5; ANL3; FHBL2
<b>Host Species</b>	Rabbit
<b>Description</b>	Anti-ANGPTL3 antibody(DM205); Rabbit mAb
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	Q9Y5C1
<b>IgG type</b>	Rabbit IgG
<b>Clonality</b>	Monoclonal
<b>Reactivity</b>	Human
<b>Applications</b>	ELISA; Flow Cyt
<b>Recommended Dilutions</b>	ELISA 1:5000-10000; Flow Cyt 1:100
<b>Purification</b>	Purified from cell culture supernatant by affinity chromatography
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	This gene encodes a member of a family of secreted proteins that function in angiogenesis. The encoded protein; which is expressed predominantly in the liver; is further processed into an N-terminal coiled-coil domain-containing chain and a C-terminal fibrinogen chain. The N-terminal chain is important for lipid metabolism; while the C-terminal chain may be involved in angiogenesis. Mutations in this gene cause familial hypobetalipoproteinemia type 2.
<b>Usage</b>	Research use only





**Figure 1.** Flow cytometry analysis with Anti-ANGPTL3 (DM205) on Expi293 cells transfected with human ANGPTL3 (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).

DIMABIO CONFIDENTIAL

