

**PRODUCT INFORMATION**

<b>Clone ID</b>	3D7
<b>Target</b>	APP
<b>Synonyms</b>	AAA;ABETA;ABPP;AD1;APPI;CTFgamma;CVAP;PN-II;PN2;preA4
<b>Host Species</b>	Rabbit
<b>Description</b>	Anti-APP antibody(3D7); IgG1 Chimeric mAb
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	P05067
<b>IgG type</b>	Rabbit/Human Fc chimeric IgG1
<b>Clonality</b>	Monoclonal
<b>Reactivity</b>	Human
<b>Applications</b>	Flow Cyt
<b>Recommended Dilutions</b>	Flow Cyt 1/100
<b>Purification</b>	Purified from cell culture supernatant by affinity chromatography
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	This gene encodes a cell surface receptor and transmembrane precursor protein that is cleaved by secretases to form a number of peptides. Some of these peptides are secreted and can bind to the acetyltransferase complex APBB1/TIP60 to promote transcriptional activation, while others form the protein basis of the amyloid plaques found in the brains of patients with Alzheimer disease. In addition, two of the peptides are antimicrobial peptides, having been shown to have bacteriocidal and antifungal activities. Mutations in this gene have been implicated in autosomal dominant Alzheimer disease and cerebroarterial amyloidosis (cerebral amyloid angiopathy). Multiple transcript variants encoding several different isoforms have been found for this gene. [provided by RefSeq, Aug 2014]
<b>Usage</b>	Research use only






Figure 1. Flow cytometry analysis with Anti-APP (3D7) mAb on Expi293 cells transfected with human APP (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).




Figure 2. Flow cytometry analysis of antigen binding of anti-human APP mAb(DMC101002).

(A) DMC101002 does not bind to CHO-S cells that do not express APP.

(B) A clear peak shift of DMC101002 was seen compared to the control when incubated with APP-expressing Siha cells, indicating strong binding of DMC101002 to APP. Antibodies were incubated at 5  $\mu\text{g}/\text{mL}$ .

