

## **PRODUCT INFORMATION**

2857916, GSK2857916, J6M0, J6M0-mcMMAF, Common Name

Unconjugated mAb

Conjugate Unconjugated

CD269;TNFRSF17;BCM;BCMA Synonyms

**Applications** ELISA; Flow Cyt

Recommended

ELISA 1:5000-10000; Flow Cyt 1:100 Dilutions

Formulation & Reconstitution Lyophilized from sterile PBS, pH 7.4. Normally 5 % 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis

for specific instructions of reconstitution.

**Host Species** Humanized

IgG type lgG1 Reactivity Human ВСМА **Target Uniprot ID** Q02223

**Description** Anti-BCMA (belantamab biosimilar) mAb

**Delivery** In Stock

Storage & Shipping

**Background** 

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store

at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

témperature.

Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans

or animals. Our unconjugated biosimilar monoclonal antibodies (mAbs) are based on the sequences outlined in relevant patents or scientific publications. These antibodies are in

their native, unconjugated form, meaning they do not contain any payload or therapeutic agent attached. They are designed for use in research and development, and their performance has been tested as standalone molecules through

comprehensive QC tests.

Usage Research use only

> All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or

**DIMA Disclaimer** reverse engineering attempt is prohibited. We are


actively scrutinizing all patent application to

Email: info@dimabio.com Website: www.dimabio.com

ensure no IP infringement.



## Anti-BCMA (belantamab biosimilar) mAb ELISA



**Figure 1.** ELISA plate pre-coated by 2  $\mu$ g/ml (100  $\mu$ l/well) Human BCMA, mFc tagged protein ([getskuurl sku="PME100035"]) can bind Anti-BCMA Neutralizing antibody in a linear range of 0.64-80.0 ng/ml



