

PRODUCT INFORMATION

Common Name	TBC
Synonyms	ABCD-2;CC chemokine TARC;C-C motif chemokine 17;CCL17;chemokine (C-C motif) ligand 17;MGC138273;SCYA17MGC138271;small inducible cytokine subfamily A (Cys-Cys);member 17;Small-inducible cytokine A17;T cell-directed CC chemokine;TARC;TARCA-152E5.3;Thymus and activation-regulated chemokine
Applications	N/A
Recommended Dilutions	N/A
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Host Species	Whole mAb
IgG type	IgG2
Reactivity	Human
Target	CCL17
Uniprot ID	Q92583
Description	Anti-CCL17(denrakibart biosimilar) IgG2) mAb
Delivery	3-4 weeks
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
Usage	Research use only
Conjugate	Unconjugated

DIMA BIOTECH CONFIDENTIAL

