

PRODUCT INFORMATION

Clone ID	DMC436
Target	CD83
Synonyms	BL11; HB15
Host Species	Rabbit
Description	Anti-CD83 antibody(DMC436); IgG1 Chimeric mAb
Delivery	In Stock
Uniprot ID	Q01151
IgG type	Rabbit/Human Fc chimeric IgG1
Clonality	Monoclonal
Reactivity	Human
Applications	Flow Cyt
Recommended Dilutions	Flow Cyt 1:100
Purification	Purified from cell culture supernatant by affinity chromatography
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The protein encoded by this gene is a single-pass type I membrane protein and member of the immunoglobulin superfamily of receptors. The encoded protein may be involved in the regulation of antigen presentation. A soluble form of this protein can bind to dendritic cells and inhibit their maturation. Three transcript variants encoding different isoforms have been found for this gene.
Usage	Research use only
Conjugate	Unconjugated
DIMA Disclaimer	All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement.






Figure 1. Flow cytometry analysis with Anti-CD83 (DMC436) on Expi293 cells transfected with human CD83 (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).




Figure 2. Flow cytometry analysis of antigen binding of anti-human CD83 mAb(DMC100436).
(A) DMC100436 does not bind to CHO-S cells that do not express CD83.
(B) A clear peak shift of DMC100436 was seen compared to the control when incubated with CD83-expressing Raji cells, indicating strong binding of DMC100436 to CD83. Antibodies were incubated at 5 µg/mL.

