

PRODUCT INFORMATION

| | |
|---|--|
| Clone ID | DM208 |
| Target | CXCR3 |
| Synonyms | CD182; CD183; CKR-L2; CMKAR3; GPR9; IP10-R; Mig-R; MigR |
| Host Species | Rabbit |
| Description | Anti-CXCR3 antibody(DM208); Rabbit mAb |
| Delivery | In Stock |
| Uniprot ID | P49682 |
| IgG type | Rabbit IgG |
| Clonality | Monoclonal |
| Reactivity | Human |
| Applications | ELISA; Flow Cyt |
| Recommended Dilutions | ELISA 1:5000-10000; Flow Cyt 1:100 |
| Purification | Purified from cell culture supernatant by affinity chromatography |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | This gene encodes a G protein-coupled receptor with selectivity for three chemokines; termed CXCL9:Mig (monokine induced by interferon-g); CXCL10:IP10 (interferon-g-inducible 10 kDa protein) and CXCL11:I-TAC (interferon-inducible T cell a-chemoattractant). Binding of chemokines to this protein induces cellular responses that are involved in leukocyte traffic; most notably integrin activation; cytoskeletal changes and chemotactic migration. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. One of the isoforms (CXCR3-B) shows high affinity binding to chemokine; CXCL4:PF4 (PMID:12782716). [provided by RefSeq; Jun 2011] |
| Usage | Research use only |






Figure 1. Flow cytometry analysis with Anti-CXCR3 (DM208) on Expi293 cells transfected with human CXCR3 (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).

DIMABIO CONFIDENTIAL

