

PRODUCT INFORMATION

| Common Name | GC33, RG7686, RO5137382 |
|---------------------------------|---|
| Synonyms | DGSX;GTR2-2;MXR7;OCI-5;SDYS;SGB;SGBS;SGBS1 |
| Conjugate | Unconjugated |
| Applications | ELISA; Flow Cyt |
| Recommended Dilutions | ELISA 1:5000-10000; Flow Cyt 1:100 |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Host Species | Humanized |
| lgG type | lgG1 |
| Reactivity | Human |
| Target | GPC3 |
| Uniprot ID | P51654 |
| Description | Anti-GPC3(codrituzumab biosimilar) mAb |
| Delivery | In Stock |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals. |
| Usage | Research use only |
| DIMA Disclaimer | All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement. |
| | |

Email: info@dimabio.com Website: www.dimabio.com





Anti-GPC3 (codrituzumab biosimilar) mAb ELISA

0.1 µg of Human GPC3, hFc tagged protein per well




Figure 1. ELISA plate pre-coated by 1 μ g/mL (100 μ L/well) Human GPC3 Protein, hFc Tag (PME100113) can bind Anti-GPC3 Neutralizing antibody (BME100083) in a linear range of 1.12-508.85 ng/mL. In order to specifically detect BME100083, mouse anti-human Fab-specific antibody was used as detection antibody.




Figure 2. Flow cytometry analysis of Huh-7 cells stained with 1 μ g/mL Anti-GPC3 (codrituzumab biosimilar) mAb (BME100083) (Blue histogram) and unstained Huh-7 cells as a negative control (Red histogram).

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com

