

PRODUCT INFORMATION

Common Name	Hen egg Lysozyme
Conjugate	Unconjugated
Synonyms	HEL
Applications	ELISA, Flow Cyt
Recommended Dilutions	ELISA 1:5000-10000, Flow Cyt 1:100
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Host Species	Chimeric
IgG type	IgG2
Reactivity	N/A
Target	HEL
Uniprot ID	N/A
Description	Anti-HEL Human IgG2-Kappa Isotype control mAb
Delivery	In Stock
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized antibodies are shipped at ambient temperature.
Background	Anti Chicken Hen Egg Lysozyme, specifically recognises Hen Egg Lysozyme (HEL), known also as muramidase or N-acetylmuramide glycanhydrolase, a 14kDa enzymic protein involved in the destruction of bacteria. Lysozyme damages bacterial cell walls by catalyzing hydrolysis of 1,4-beta-linkages between N-acetylmuramic acid and N-acetyl-D-glucosamine residues in a peptidoglycan and between N-acetyl-D-glucosamine residues in chitodextrins. Lysozyme is abundant in a number of secretions, such as tears, saliva, human milk and mucus. It is also present in cytoplasmic granules of PMN's and high concentrations of lysozyme are present in egg white. C-type lysozymes are closely related to alpha-lactalbumin in sequence and structure making them part of the same family.
Usage	Research use only
DIMA Disclaimer	All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement.

