

PRODUCT INFORMATION

4D5-8,4D5V8, Herceptin, rhuMabHER2, Unconjugated **Common Name**

mAb

Synonyms ERBB2;CD340;HER-2/neu;HER2;MLN19;NEU;NGL;TKR1

Conjugate Unconjugated ELISA; Flow Cyt **Applications**

Recommended

ELISA 1:5000-10000; Flow Cyt 1:100 **Dilutions**

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for Formulation & Reconstitution

specific instructions of reconstitution.

Host Species Humanized

IgG type laG1 Reactivity Human **Target** Her2 P04626 **Uniprot ID**

Description Anti-Her2 (trastuzumab biosimilar)mAb

Delivery In Stock

Storage & Shipping

Background

DIMA Disclaimer

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid

repeated freezing and thawing). Lyophilized proteins

are shipped at ambient temperature

Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals. Our unconjugated biosimilar monoclonal antibodies (mAbs) are based on the sequences outlined in relevant patents or scientific publications. These antibodies are in their native, unconjugated form, meaning they do not contain any payload or therapeutic agent attached. They are designed for

use in research and development, and their performance has been tested as standalone molecules through comprehensive QC tests.

Usage Research use only

All DIMA recombinant antibodies are genuinely

generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP

Email: info@dimabio.com Website: www.dimabio.com

infringement.

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China)

/+86-400-006-0995(China)





Anti-Her2 (trastuzumab biosimilar) mAb ELISA

0.1 µg of Human Her2, His tagged protein per well




Figure 1. ELISA plate pre-coated by 1 μ g/mL (100 μ L/well) Human Her2, His tagged protein (PME100095) can bind Anti-Her2 (trastuzumab biosimilar) mAb (BME100048) in a linear range of 3.2-400 ng/ml.

Anti-Her2 (trastuzumab biosimilar)mAb ELISA

0.1 μg of Human Her2, hFc tagged protein per well




Figure 2. ELISA plate pre-coated by 1 μ g/mL (100 μ L/well) Human Her2 Protein, hFc Tag (PME100665) can bind Anti-Her2 (trastuzumab biosimilar) mAb (BME100048) in a linear range of 3.81-1730.10 ng/mL. In order to specifically detect BME100048, mouse anti-human Fab-specific antibody was used as detection antibody.

Email: info@dimabio.com Website: www.dimabio.com








Figure 3. Flow cytometry analysis with 15 μ g/mL Anti-Her2 (trastuzumab biosimilar) mAb (BME100048) on Expi293 cells transfected with Human Her2 protein (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).





Figure 4. Flow cytometry analysis of antigen binding of anti-human Her2 mAb(BME100048). (A) BME100048 does not bind to Jurkat cells that do not express Her2. (B) A clear peak shift of BME100048 was seen compared to the control when incubated with Her2-expressing MCF-7 cells, indicating strong binding of BME100048 to Her2. Antibodies were incubated at 2 μ g/mL.



Email: info@dimabio.com Website: www.dimabio.com

