


**PRODUCT INFORMATION**

<b>Clone ID</b>	DM183
<b>Target</b>	IL13RA1
<b>Synonyms</b>	CD213A1; CT19; IL-13Ra; NR4
<b>Host Species</b>	Rabbit
<b>Description</b>	Anti-IL13RA1 antibody(DM183); Rabbit mAb
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	P78552
<b>IgG type</b>	Rabbit IgG
<b>Clonality</b>	Monoclonal
<b>Reactivity</b>	Human
<b>Applications</b>	ELISA; Flow Cyt
<b>Recommended Dilutions</b>	ELISA 1:5000-10000; Flow Cyt 1:100
<b>Purification</b>	Purified from cell culture supernatant by affinity chromatography
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	The protein encoded by this gene is a subunit of the interleukin 13 receptor. This subunit forms a receptor complex with IL4 receptor alpha; a subunit shared by IL13 and IL4 receptors. This subunit serves as a primary IL13-binding subunit of the IL13 receptor; and may also be a component of IL4 receptors. This protein has been shown to bind tyrosine kinase TYK2; and thus may mediate the signaling processes that lead to the activation of JAK1; STAT3 and STAT6 induced by IL13 and IL4.
<b>Usage</b>	Research use only





**Figure 1.** Flow cytometry analysis with Anti-IL13RA1 (DM183) on Expi293 cells transfected with human IL13RA1 (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).

DIMABIO CONFIDENTIAL

