

PRODUCT INFORMATION

Clone ID	1G1
Target	IL18BP
Synonyms	FVH;IL18BPa
Host Species	Rabbit
Description	Anti-IL18BP antibody(1G1), IgG1 Chimeric mAb
Delivery	In Stock
Uniprot ID	O95998
lgG type	Rabbit/Human Fc chimeric IgG1
Clonality	Monoclonal
Reactivity	Human
Applications	ELISA
Recommended Dilutions	ELISA 1/5000-10000
Purification	Purified from cell culture supernatant by affinity chromatography
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The protein encoded by this gene functions as an inhibitor of the proinflammatory cytokine, IL18. It binds IL18, prevents the binding of IL18 to its receptor, and thus inhibits IL18-induced IFN- gamma production, resulting in reduced T-helper type 1 immune responses. This protein is constitutively expressed and secreted in mononuclear cells. Elevated level of this protein is detected in the intestinal tissues of patients with Crohn's disease. Alternatively spliced transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Feb 2011]
Usage	Research use only
Conjugate	Unconjugated
DIMA Disclaimer	All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement.

Email: info@dimabio.com Website: www.dimabio.com








Figure 1. ELISA plate pre-coated by 1 μ g/ml (100 μ l/well) Human IL18BP, hFc tagged protein can bind Rabbit anti-IL18BP monoclonal antibody(clone: 1G1) in a linear range of 5-100 ng/ml.

Email: info@dimabio.com Website: www.dimabio.com

