

PRODUCT INFORMATION

Clone ID	DMC393
Target	IL5RA
Synonyms	IL5Ra;CD125;IL-5 R alpha
Host Species	Rabbit
Description	Anti-IL5RA antibody(DMC393); IgG1 Chimeric mAb
Delivery	In Stock
Uniprot ID	Q01344
IgG type	Rabbit/Human Fc chimeric IgG1
Clonality	Monoclonal
Reactivity	Human
Applications	Flow Cyt
Recommended Dilutions	Flow Cyt 1:100
Purification	Purified from cell culture supernatant by affinity chromatography
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The protein encoded by this gene is an interleukin 5 specific subunit of a heterodimeric cytokine receptor. The receptor is comprised of a ligand specific alpha subunit and a signal transducing beta subunit shared by the receptors for interleukin 3 (IL3); colony stimulating factor 2 (CSF2:GM-CSF); and interleukin 5 (IL5). The binding of this protein to IL5 depends on the beta subunit. The beta subunit is activated by the ligand binding; and is required for the biological activities of IL5. This protein has been found to interact with syndecan binding protein (syntenin); which is required for IL5 mediated activation of the transcription factor SOX4. Several alternatively spliced transcript variants encoding four distinct isoforms have been reported.
Usage	Research use only






Figure 1. Flow cytometry analysis with Anti-IL5RA (DMC393) on Expi293 cells transfected with human IL5RA (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).

