

PRODUCT INFORMATION

| | |
|---|---|
| Common Name | IMC RON-8, IMC-RON8, RON8 |
| Synonyms | CD136;CDw136;NPCA3;PTK8;RON |
| Applications | ELISA |
| Recommended Dilutions | ELISA 1:5000-10000 |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Host Species | Homo sapiens |
| IgG type | IgG1 |
| Reactivity | Human |
| Target | MST1R |
| Uniprot ID | Q04912 |
| Description | Anti-MST1R(narnatumab biosimilar) mAb |
| Delivery | In Stock |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals. |
| Usage | Research use only |
| Conjugate | Unconjugated |

DIMA BIO CONFIDENTIAL




Anti-MST1R (narnatumab biosimilar) mAb ELISA0.2 μ g of Human MST1R, hFc tagged protein per well

Figure 1. ELISA plate pre-coated by 2 μ g/mL (100 μ L/well) Human MST1R Protein, hFc Tag (PME100821) can bind Anti-MST1R Neutralizing antibody (BME100118) in a linear range of 0.73–46.88 ng/mL. In order to specifically detect BME100118, mouse anti-human Fab-specific antibody was used as detection antibody.

