

**PRODUCT INFORMATION**

<b>Clone ID</b>	DMC683
<b>Target</b>	TENM4
<b>Synonyms</b>	Doc4; ETM5; ODZ4; ten-4; Ten-M4; TEN4; TNM4
<b>Host Species</b>	Rabbit
<b>Description</b>	Anti-TENM4 antibody(DMC683); IgG1 Chimeric mAb
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	Q6N022
<b>IgG type</b>	Rabbit/Human Fc chimeric IgG1
<b>Clonality</b>	Monoclonal
<b>Reactivity</b>	Human
<b>Applications</b>	Flow Cyt
<b>Recommended Dilutions</b>	Flow Cyt 1:100
<b>Purification</b>	Purified from cell culture supernatant by affinity chromatography
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	The protein encoded by this gene plays a role in establishing proper neuronal connectivity during development. Defects in this gene have been associated with hereditary essential tremor-5. [provided by RefSeq; Oct 2016]
<b>Usage</b>	Research use only






Figure 1. Flow cytometry analysis with Anti-TENM4(DMC683) on Expi293 cells transfected with human TENM4(Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).




Figure 2. Flow cytometry analysis of antigen binding of anti-human TENM4 mAb(DMC100683).

(A) DMC100683 does not bind to CHO-S cells that do not express TENM4.

(B) A clear peak shift of DMC100683 was seen compared to the control when incubated with TENM4-expressing TT cells, indicating strong binding of DMC100683 to TENM4. Antibodies were incubated at 5  $\mu$ g/mL.

