

## **PRODUCT INFORMATION**

Clone ID **DMC477 Target** CCR6

BN-1; C-C CKR-6; CC-CKR-6; CCR-6; CD196; CKR-**Synonyms** 

L3; CKRL3; CMKBR6; DCR2; DRY6; GPR29;

GPRCY4; STRL22

**Host Species** 

Biotinylated Anti-CCR6 antibody(DMC477); IgG1 **Description** 

Chimeric mAb

Delivery 2-3 weeks P51684 **Uniprot ID** 

Rabbit/Human Fc chimeric IgG1 IgG type

Monoclonal Clonality Reactivity Human **Applications** Flow Cyt

Recommended

Storage & Shipping

**Background** 

Flow Cyt 1:100 **Dilutions** 

Purified from cell culture supernatant by affinity **Purification** 

chromatography

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before Formulation & lyophilization. Please see Certificate of Analysis Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

temperature.

This gene encodes a member of the beta chemokine receptor family; which is predicted to

be a seven transmembrane protein similar to G protein-coupled receptors. The gene is preferentially expressed by immature dendritic cells and memory T cells. The ligand of this receptor is macrophage inflammatory protein 3 alpha (MIP-3 alpha). This receptor has been the beautiful to the protein to the ligand maturation. shown to be important for B-lineage maturation and antigen-driven B-cell differentiation; and it

may regulate the migration and recruitment of dentritic and T cells during inflammatory and immunological responses. Alternatively spliced transcript variants that encode the same protein have been described for this gene. [provided by

> Email: info@dimabio.com Website: www.dimabio.com

RefSeq; Jul 2008]

**Usage** Research use only

Conjugate Biotinylated

