Cat. No. DMC100270B



## **PRODUCT INFORMATION**

Clone ID DMC270
Target CD160

**Synonyms** BY55; NK1; NK28

**Host Species** Rabbit

**Description**Biotinylated Anti-CD160 antibody(DMC270); IgG1

Chimeric mAb

**Delivery** 2-3 weeks **Uniprot ID** 095971

**IgG type** Rabbit/Human Fc chimeric IgG1

Clonality Monoclonal
Reactivity Human
Applications Flow Cyt

Recommended

Dilutions

Storage & Shipping

Background

Flow Cyt 1:100

**Purification**Purified from cell culture supernatant by affinity

chromatography

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis

lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

témperature.

CD160 is an 27 kDa glycoprotein which was initially identified with the monoclonal antibody BY55. Its expression is tightly associated with peripheral blood NK cells and CD8 T lymphocytes with cytolytic effector activity. The cDNA sequence of CD160 predicts a cysteine-rich;

with cytolytic effector activity. The cDNA sequence of CD160 predicts a cysteine-rich; glycosylphosphatidylinositol-anchored protein of 181 amino acids with a single Ig-like domain weakly homologous to KIR2DL4 molecule. CD160 is expressed at the cell surface as a tightly disulfide-linked multimer. RNA blot analysis

revealed CD160 mRNAs of 1.5 and 1.6 kb whose expression was highly restricted to circulating NK and T cells; spleen and small intestine. Within NK cells CD160 is expressed by CD56dimCD16 cells whereas among circulating T cells its expression is mainly restricted to TCRgd bearing cells and to TCRab CD8brightCD95 CD56 CD28-CD27-cells. In tissues; CD160 is expressed on all intestinal intraepithelial lymphocytes. CD160 shows a broad specificity for binding to both classical and

nonclassical MHC class I molecules.

**Usage** Research use only

**Conjugate** Biotinylated

All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are

actively scrutinizing all patent application to

ensure no IP infringement.

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China)

**DIMA Disclaimer** 

/+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com



Biotinylated Anti-CD160 antibody(DMC270); IgG1 Chimeric mAb

Email: info@dimabio.com Website: www.dimabio.com

Cat. No. DMC100270B





