

PRODUCT INFORMATION

|                              |   |
|------------------------------|---|
| Clone ID                     | DM9   |
| Target                       | CS1   |
| Synonyms                     | SLAM7 (19A; CD319; CRACC; CS1)  |
| Host Species                 | Rabbit  |
| Description                  | Biotinylated Anti-CS1 antibody(DM9); Rabbit mAb   |
| Delivery                     | 2-3 weeks   |
| Uniprot ID                   | Q9NQ25  |
| IgG type                     | Rabbit IgG  |
| Clonality                    | Monoclonal  |
| Reactivity                   | Human   |
| Applications                 | ELISA; Flow Cyt   |
| Recommended Dilutions        | Flow Cyt 1:100  |
| Purification                 | Purified from cell culture supernatant by affinity chromatography   |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.  |
| Storage & Shipping           | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.   |
| Background                   | Self-ligand receptor of the signaling lymphocytic activation molecule (SLAM) family. SLAM receptors triggered by homo- or heterotypic cell-cell interactions are modulating the activation and differentiation of a wide variety of immune cells and thus are involved in the regulation and interconnection of both innate and adaptive immune response. Activities are controlled by presence or absence of small cytoplasmic adapter proteins; SH2D1A:SAP and/or SH2D1B:EAT-2. Isoform 1 mediates NK cell activation through a SH2D1A-independent extracellular signal-regulated ERK-mediated pathway (PubMed:11698418). Positively regulates NK cell functions by a mechanism dependent on phosphorylated SH2D1B. Downstream signaling implicates PLCG1; PLCG2 and PI3K (PubMed:16339536). In addition to heterotypic NK cells-target cells interactions also homotypic interactions between NK cells may contribute to activation. However; in the absence of SH2D1B; inhibits NK cell function. Acts also inhibitory in T-cells (By similarity). May play a role in lymphocyte adhesion (PubMed:11802771). In LPS-activated monocytes negatively regulates production of proinflammatory cytokines (PubMed:23695528). |
| Usage                        | Research use only   |
| Conjugate                    | Biotinylated  |



**DIMA Disclaimer**

All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement.

DIMABIO CONFIDENTIAL

