

PRODUCT INFORMATION

Clone ID	DMC500
Target	HBEGF
Synonyms	DTR; DTS; DTSF; HEGFL
Host Species	Rabbit
Description	Biotinylated Anti-HBEGF antibody(DMC500); IgG1 Chimeric mAb
Delivery	2-3 weeks
Uniprot ID	Q99075
IgG type	Rabbit/Human Fc chimeric IgG1
Clonality	Monoclonal
Reactivity	Human
Applications	Flow Cyt
Recommended Dilutions	Flow Cyt 1:100
Purification	Purified from cell culture supernatant by affinity chromatography
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Growth factor that mediates its effects via EGFR; ERBB2 and ERBB4. Required for normal cardiac valve formation and normal heart function. Promotes smooth muscle cell proliferation. May be involved in macrophage-mediated cellular proliferation. It is mitogenic for fibroblasts; but not endothelial cells. It is able to bind EGF receptor:EGFR with higher affinity than EGF itself and is a far more potent mitogen for smooth muscle cells than EGF. Also acts as a diphtheria toxin receptor.[UniProtKB:Swiss-Prot Function]
Usage	Research use only
Conjugate	Biotinylated

