

**PRODUCT INFORMATION**

<b>Clone ID</b>	DM168
<b>Target</b>	MICB
<b>Synonyms</b>	MIC-B; PERB11.2
<b>Host Species</b>	Rabbit
<b>Description</b>	Anti-MICB antibody(DM168); Rabbit mAb
<b>Delivery</b>	2-3 weeks
<b>Uniprot ID</b>	Q29980
<b>IgG type</b>	Rabbit IgG
<b>Clonality</b>	Monoclonal
<b>Reactivity</b>	Human
<b>Applications</b>	ELISA; Flow Cyt
<b>Recommended Dilutions</b>	ELISA 1:5000-10000; Flow Cyt 1:100
<b>Purification</b>	Purified from cell culture supernatant by affinity chromatography
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	This gene encodes a heavily glycosylated protein which is a ligand for the NKG2D type II receptor. Binding of the ligand activates the cytolytic response of natural killer (NK) cells; CD8 alphabeta T cells; and gammadelta T cells which express the receptor. This protein is stress-induced and is similar to MHC class I molecules; however; it does not associate with beta-2-microglobulin or bind peptides. Alternative splicing results in multiple transcript variants.
<b>Usage</b>	Research use only

