

PRODUCT INFORMATION

Target MMAE
Synonyms N.A.
Host Species Rabbit

DescriptionBiotinylated Anti-MMAE antibody(11B2); Rabbit

mAb

Uniprot ID N.A.

IgG type Rabbit IgG
Clonality Monoclonal

Reactivity N.A. **Applications** ELISA

Recommended Dilutions ELISA 1:5000-10000

Delivery In Stock

PurificationPurified from cell culture supernatant by affinity

chromatography

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

Storage & Shipping intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

Email: info@dimabio.com Website: www.dimabio.com

témperature.

Background N.A

Usage Research use only

Conjugate Biotinylated





Elisa assay to evaluate Biotinylated Anti-MMAE antibody 0.2μg Human IgG MMAE per well




Figure 1. Elisa plates were pre-coated with IgG-MMAE ($0.2\mu g/per$ well). Serial diluted Biotinylated Anti-MMAE monoclonal antibody (DME101006B) solutions were added, washed, and incubated with SA-HRP before Elisa reading. From above data, the EC50 for anti-MMAE monoclonal antibody binding with IgG-MMAE is 9.72 ng/ml.

Email: info@dimabio.com Website: www.dimabio.com

