

**PRODUCT INFORMATION**

<b>Clone ID</b>	DM115
<b>Target</b>	TNFRSF10B
<b>Synonyms</b>	TNFRSF10B;TRAILR2;TRAIL-R2;CD262;DR5;KILLER;TRICK2;ZTNFR9;TRICKB
<b>Host Species</b>	Rabbit
<b>Description</b>	Anti-TNFRSF10B antibody(DM115); Rabbit mAb
<b>Delivery</b>	2-3 weeks
<b>Uniprot ID</b>	O14763
<b>IgG type</b>	Rabbit IgG
<b>Clonality</b>	Monoclonal
<b>Reactivity</b>	Human
<b>Applications</b>	ELISA; Flow Cyt
<b>Recommended Dilutions</b>	ELISA 1:5000-10000; Flow Cyt 1:100
<b>Purification</b>	Purified from cell culture supernatant by affinity chromatography
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	The protein encoded by this gene is a member of the TNF-receptor superfamily; and contains an intracellular death domain. This receptor can be activated by tumor necrosis factor-related apoptosis inducing ligand (TNFSF10:TRAIL:APO-2L); and transduces an apoptosis signal. Studies with FADD-deficient mice suggested that FADD; a death domain containing adaptor protein; is required for the apoptosis mediated by this protein. Two transcript variants encoding different isoforms and one non-coding transcript have been found for this gene.
<b>Usage</b>	Research use only

