

**PRODUCT INFORMATION**

|   |   |
|---|---|
| <b>Target</b>                           | ADAM9   |
| <b>Synonyms</b>                         | CORD9;MCMP;MDC9;Mltng   |
| <b>Description</b>                      | Recombinant Cynomolgus ADAM9 protein with C-terminal 6×His tag  |
| <b>Delivery</b>                         | In Stock  |
| <b>Uniprot ID</b>                       | A0A2K5X4X8  |
| <b>Expression Host</b>                  | HEK293  |
| <b>Tag</b>                              | C-6×His Tag   |
| <b>Molecular Characterization</b>       | ADAM9(Ala29-Gly698) 6×His tag   |
| <b>Molecular Weight</b>                 | The protein has a predicted molecular mass of 74.9 kDa after removal of the signal peptide. The apparent molecular mass of cADAM9-His is approximately 55-100 kDa due to glycosylation.   |
| <b>Purity</b>                           | The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue staining.  |
| <b>Formulation &amp; Reconstitution</b> | Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.   |
| <b>Storage &amp; Shipping</b>           | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.   |
| <b>Background</b>                       | This gene encodes a member of the ADAM (a disintegrin and metalloprotease domain) family. Members of this family are membrane-anchored proteins structurally related to snake venom disintegrins, and have been implicated in a variety of biological processes involving cell-cell and cell-matrix interactions, including fertilization, muscle development, and neurogenesis. The protein encoded by this gene interacts with SH3 domain-containing proteins, binds mitotic arrest deficient 2 beta protein, and is also involved in TPA-induced ectodomain shedding of membrane-anchored heparin-binding EGF-like growth factor. Several alternatively spliced transcript variants have been identified for this gene. [provided by RefSeq, Jul 2010] |
| <b>Usage</b>                            | Research use only   |
| <b>Conjugate</b>                        | Unconjugated  |






Figure 1. Cynomolgus ADAM9 Protein, His Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

