

PRODUCT INFORMATION

| | |
|---|--|
| Target | CLEC4C |
| Synonyms | BDCA-2;BDCA2;CD303;CLECSF7;CLECSF11;DLEC;HECL;PRO34150 |
| Description | Recombinant Cynomolgus CLEC4C protein with N-terminal 6×His tag |
| Delivery | In Stock |
| Uniprot ID | A0A2K5UWP4 |
| Expression Host | HEK293 |
| Tag | N-6×His Tag |
| Molecular Characterization | 6×His tag CLEC4C(Tyr48-Ile212) |
| Molecular Weight | The protein has a predicted molecular mass of 20.3 kDa after removal of the signal peptide. The apparent molecular mass of His-cCLEC4C is approximately 15-25 kDa due to glycosylation. |
| Purity | The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue staining. |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | This gene encodes a member of the C-type lectin/C-type lectin-like domain (CTL/CTLD) superfamily. Members of this family share a common protein fold and have diverse functions, such as cell adhesion, cell-cell signalling, glycoprotein turnover, and roles in inflammation and immune response. The encoded type 2 transmembrane protein may play a role in dendritic cell function. Two transcript variants encoding distinct isoforms have been identified for this gene. [provided by RefSeq, Jul 2008] |
| Usage | Research use only |
| Conjugate | Unconjugated |





Figure 1. Cynomolgus CLEC4C Protein, His Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

