

PRODUCT INFORMATION

| | |
|---|---|
| Target | Her2 |
| Synonyms | NEU; NGL; ERBB2; TKR1; CD340; HER-2; VSCN2; MLN 19; MLN-19; c-ERB2; c-ERB-2; HER-2/neu; p185(erbB2) |
| Description | Recombinant Cynomolgus Her2 protein with C-terminal 10×His tag |
| Delivery | In Stock |
| Uniprot ID | XP_005584091.3 |
| Expression Host | HEK293 |
| Tag | C-10×His tag |
| Molecular Characterization | Her2(Thr120-Thr749) 10×His tag |
| Molecular Weight | The protein has a predicted molecular mass of 70.7 kDa after removal of the signal peptide. The apparent molecular mass of cHer2-His is approximately 70-130 kDa due to glycosylation. |
| Purity | The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue staining. |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | This gene encodes a member of the epidermal growth factor (EGF) receptor family of receptor tyrosine kinases. This protein has no ligand binding domain of its own and therefore cannot bind growth factors. However, it does bind tightly to other ligand-bound EGF receptor family members to form a heterodimer, stabilizing ligand binding and enhancing kinase-mediated activation of downstream signalling pathways, such as those involving mitogen-activated protein kinase and phosphatidylinositol-3 kinase. Allelic variations at amino acid positions 654 and 655 of isoform a (positions 624 and 625 of isoform b) have been reported, with the most common allele, Ile654/Ile655, shown here. Amplification and/or overexpression of this gene has been reported in numerous cancers, including breast and ovarian tumors. Alternative splicing results in several additional transcript variants, some encoding different isoforms and others that have not been fully characterized. [provided by RefSeq, Jul 2008] |
| Usage | Research use only |
| Conjugate | Unconjugated |






Figure 1. Cynomolgus Her2 Protein, His Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

