

## **PRODUCT INFORMATION**

HER3 **Target** 

ERBB3; FERLK; LCCS2; VSCN1; ErbB-3; c-erbB3; **Synonyms** erbB3-S; MDA-BF-1; c-erbB-3; p180-ErbB3; p45-

sErbB3; p85-sErbB3

Recombinant Cynomolgus HER3 protein with C-**Description** 

terminal 10×His tag

Delivery In Stock **Uniprot ID** G7PIG5 **Expression Host HEK293** 

Tag C-10×His tag

Molecular Characterization

Background

HER3(Ser20-Thr643) 10×His tag

The protein has a predicted molecular mass of 70.0 kDa after removal of the signal peptide. The apparent molecular mass of cHER3-His is **Molecular Weight** 

approximately 100-130 kDa due to glycosylation. The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue

**Purity** 

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before Formulation & Reconstitution lyophilization. Please see Certificate of Analysis

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

témperature.

This gene encodes a member of the epidermal growth factor receptor (EGFR) family of receptor tyrosine kinases. This membrane-bound protein has a neuregulin binding domain but not an active kinase domain. It therefore can bind this ligand but not convey the signal into the cell through protein phosphorylation. However, it does form heterodimers with other EGF receptor family members which do have kinase activity. Heterodimerization leads to the activation of pathways which lead to cell proliferation or differentiation. Amplification of this gene and/or overexpression of its protein have been reported

in numerous cancers, including prostate, bladder, and breast tumors. Alternate transcriptional splice variants encoding different isoforms have been

characterized. One isoform lacks the

intermembrane region and is secreted outside the cell. This form acts to modulate the activity of the membrane-bound form. Additional splice variants have also been reported, but they have not been thoroughly characterized. [provided by RefSeq, Jul

> Email: info@dimabio.com Website: www.dimabio.com

20081

Usage Research use only Conjugate Unconjugated

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)






Figure 1. Cynomolgus HER3 Protein, His Tag on SDS-PAGE under reducing condition.

