Warranty and Disclaimer

Background



PRODUCT INFORMATION

CCR5 **Target**

Monoclonal Cell Line Derived from K562 Cells, Engineered for Stable Expression of Human CCR5 Using Lentiviral Technology **Description**

Host Cells **Uniprot ID** P51681 **FACS Data Applications**

RPMI-1640+10% FBS+1% P.S+1% Gln+2 ug/mL Puromycin **Growth media**

5F6 Cells/ml Package **Host Species** Human

Suggested Control SKU: BME100204

> 1. Please inspect cells upon receipt and report any issues promptly. 2. We offer one-time replacements for issues reported within a week of receipt. 3. User-induced issues are not eligible for free replacements. 4. We do not accept liability for damages resulting from cell use, storage, or loss. 5.

> Feedback received more than one month after receipt will not be processed.

Cells are shipped using dry ice and require liquid nitrogen storage for long term preservation. Storage & Shipping

Synonyms CC-CKR-5;CCCKR5;CCR-5;CD195;CKR-5;CKR5;CMKBR5;IDDM22

> This gene encodes a member of the beta chemokine receptor family, which is predicted to be a seven transmembrane protein similar to G protein-coupled receptors. This protein is expressed by T cells and macrophages, and is known to be an important co-receptor for macrophage-tropic virus, including HIV, to enter host cells. Defective alleles of this gene have been associated with the HIV infection resistance. The ligands of this receptor include monocyte chemoattractant protein 2

> of this receptor include monocyte chemoattractant protein 2 (MCP-2), macrophage inflammatory protein 1 alpha (MIP-1 alpha), macrophage inflammatory protein 1 beta (MIP-1 beta) and regulated on activation normal T expressed and secreted protein (RANTES). Expression of this gene was also detected in a promyeloblastic cell line, suggesting that this protein may play a role in granulocyte lineage proliferation and differentiation. This gene is located at the chemokine receptor gene cluster region. An allelic polymorphism in this gene results in both functional and non-functional alleles; the reference genome represents the functional allele. Two transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Jul 2015]

Email: info@dimabio.com Website: www.dimabio.com

Usage For research use only.

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)








Figure 1. Flow cytometry analysis of human CCR5 overexpression using Hu_CCR5 K562 Cell Line (Cat. No. CEL100090) and Anti-CCR5(leronlimab biosimilar) mAb (Cat. No. BME100204)

Email: info@dimabio.com Website: www.dimabio.com

