

PRODUCT INFORMATION

NTB-A **Target**

Monoclonal Cell Line Derived from CHO-S Cells, Description Engineered for Stable Expression of Human NTB-A

Using Lentiviral Technology

Host Cells CHO-S Q96DU3 **Uniprot ID FACS Data** Applications

DMEM+10% FBS+1% P.S+Gln+2 ug/mL **Growth media**

Puromycin **Package** 5E6 Cells/mL

Host Species Human

Warranty and

Disclaimer

Background

SKU: DME100159 **Suggested Control**

1. Please inspect cells upon receipt and report any issues promptly. 2. We offer one-time replacements for issues reported within a week of receipt. 3. User-induced issues are not eligible for free replacements. 4. We do not accept liability for damages resulting from cell use, storage, or loss. 5. Feedback received more than one month

after receipt will not be processed.

Cells are shipped using dry ice and require liquid Storage & Shipping nitrogen storage for long term preservation.

Synonyms NTB-A;SLAMF6;Ly108;NK-T-B-antigen;CD352;KALI


> The protein encoded by this gene is a type I transmembrane protein; belonging to the CD2 subfamily of the immunoglobulin superfamily. This encoded protein is expressed on Natural killer (NK); T; and B lymphocytes. It undergoes tyrosine phosphorylation and associates with the Src homology 2 domain-containing protein (SH2D1A) as well as with SH2 domain-containing

> phosphatases (SHPs). It functions as a coreceptor in the process of NK cell activation. It can also mediate inhibitory signals in NK cells from Xlinked lymphoproliferative patients. Alternative splicing results in multiple transcript variants

> > Email: info@dimabio.com Website: www.dimabio.com

encoding distinct isoforms.

Usage For research use only.







Hu_NTB-A CHO-S Cell Line




Figure 1. Flow cytometry analysis of human NTB-A overexpression using Hu_NTB-A CHO-S Cell Line (Cat. No. CEL100063) and Anti-NTB-A antibody(DM159)Rabbit mAb (Cat. No. DME100159)

Email: info@dimabio.com Website: www.dimabio.com

