

PRODUCT INFORMATION

Target	2B4
Synonyms	CD244;2B4;SLAMF4;NKR2B4;NAIL;h2B4
Description	Recombinant human 2B4 protein with C-terminal human Fc tag
Delivery	In Stock
Uniprot ID	Q9BZW8
Expression Host	HEK293
Tag	C-Human Fc Tag
Molecular Characterization	2B4(Cys22-Ala216) hFc(Glu99-Ala330)
Molecular Weight	The protein has a predicted molecular mass of 47.7 kDa after removal of the signal peptide.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene encodes a cell surface receptor expressed on natural killer (NK) cells (and some T cells) that mediate non-major histocompatibility complex (MHC) restricted killing. The interaction between NK-cell and target cells via this receptor is thought to modulate NK-cell cytolytic activity. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Human 2B4, hFc Tag on SDS-PAGE under reducing condition.

Human 2B4, hFc Tagged protein ELISA

0.2 μ g of 2B4, hFc Tagged protein per well




Figure 2. ELISA plate pre-coated by 2 μ g/ml (100 μ l/well) Human CD48, mFc-His tagged protein PME100016 can bind Human 2B4, hFc tagged protein (PME100477) in a linear range of 62.5-4000 ng/ml.

