

PRODUCT INFORMATION

| | |
|---|---|
| Tag | C-Flag Tag |
| Target | ACHA3 |
| Synonyms | BAIPRCK, LNCR2, NACHRA3, PAOD2 |
| Description | Human ACHA3 full length protein-synthetic nanodisc |
| Delivery | 6~8weeks |
| Uniprot ID | P32297 |
| Expression Host | HEK293 |
| Protein Families | Ion Channels: Cys-loop Receptors |
| Protein Pathways | N/A |
| Molecular Weight | The human full length ACHA3 protein has a MW of 57.5kDa |
| Formulation & Reconstitution | Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | This locus encodes a member of the nicotinic acetylcholine receptor family of proteins. Members of this family of proteins form pentameric complexes comprised of both alpha and beta subunits. This locus encodes an alpha-type subunit, as it contains characteristic adjacent cysteine residues. The encoded protein is a ligand-gated ion channel that likely plays a role in neurotransmission. Polymorphisms in this gene have been associated with an increased risk of smoking initiation and an increased susceptibility to lung cancer. Alternatively spliced transcript variants have been described. [provided by RefSeq, Nov 2009] |
| Usage | Research use only |
| Conjugate | Unconjugated |

