

PRODUCT INFORMATION

Tag C-Flag Tag

Target ACHA6

Synonyms CHNRA6

DescriptionHuman ACHA6 full length protein-synthetic

nanodisc

Delivery 6~8weeks

Uniprot ID Q15825

Expression Host HEK293

Protein Families Ion Channels: Cys-loop Receptors

Protein Pathways N/A

Storage & Shipping

Background

Molecular Weight

The human full length ACHA6 protein has a MW of 56 OkDa

56.9kDa Lyophilized from nanodisc solubilization buffer (20

mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5%
– 8% trehalose is added as protectants before
lyophilization. Please see Certificate of Analysis
for specific instructions. Do not use solvents with

Reconstitutionfor specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

témperature.

This gene encodes an alpha subunit of neuronal nicotinic acetylcholine receptors. These receptors consist of five subunits and function as ion channels involved in neurotransmission. The encoded protein is a subunit of neuronal nicotinic

encoded protein is a subunit of neurona acetylcholine receptors that mediate

dopaminergic neurotransmission and are activated by acetylcholine and exogenous nicotine. Alternatively spliced transcript variants have been observed for this gene. Single nucleotide polymorphisms in this gene have been associated with both nicotine and alcohol

Email: info@dimabio.com Website: www.dimabio.com

dependence. [provided by RefSeq, Dec 2010]

Usage Research use only

Conjugate Unconjugated



