

PRODUCT INFORMATION

Tag C-Flag&Strep Tag

Target ACHB

Synonyms ACHRB, CHRNB, CMS1D, CMS2A, CMS2C, SCCMS
Human ACHB-Strep full length protein-synthetic

Description Human A nanodisc

Delivery 6~8weeks
Uniprot ID P11230
Expression Host HEK293

Protein Families Ion Channels: Cys-loop Receptors

Protein Pathways N/A

Storage & Shipping

Background

Molecular Weight

The human full length ACHB-Strep protein has a MW of 56.7 kDa

MW of 56.7 kDa Lyophilized from nanodisc solubilization buffer (20

mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5%
– 8% trehalose is added as protectants before
lyophilization. Please see Certificate of Analysis
for specific instructions. Do not use solvents with

a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

témperature.

The muscle acetylcholine receptor is composed of five subunits: two alpha subunits and one beta, one gamma, and one delta subunit. This gene encodes the beta subunit of the acetylcholine receptor. The acetylcholine receptor changes conformation upon acetylcholine binding leading to the opening of an ion-conducting channel across the places a membrane. Mutations in this

to the opening of an ion-conducting channel across the plasma membrane. Mutations in this gene are associated with slow-channel congenital myasthenic syndrome. [provided by RefSeq, Jul

2008]

Usage Research use only
Conjugate Unconjugated

Email: info@dimabio.com Website: www.dimabio.com

