

**PRODUCT INFORMATION**

<b>Target</b>	ACHB4
<b>Synonyms</b>	N/A
<b>Description</b>	Human ACHB4 full length protein-synthetic nanodisc
<b>Delivery</b>	6~8weeks
<b>Uniprot ID</b>	P30926
<b>Expression Host</b>	HEK293
<b>Protein Families</b>	Ion Channels: Cys-loop Receptors
<b>Protein Pathways</b>	N/A
<b>Molecular Weight</b>	The human full length ACHB4 protein has a MW of 56.4kDa
<b>Formulation &amp; Reconstitution</b>	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with pH lower than 6.5 in subsequent experiments.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	This gene is found within a conserved gene cluster and encodes one of the beta subunits of the nicotinic acetylcholine receptor (nAChRs) superfamily which form ligand-gated ion channels with a central pore that forms a cation channel. Neuronal nAChRs are pentameric structures that can be either homomeric or heteromeric, with heteromeric structures containing both alpha and beta subunits. Each subunit contains an extracellular amino terminus and four transmembrane domains. Nicotine is one of the agonists that binds to the receptor. Variants in this gene have been associated with nicotine dependence and lung cancer. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Sep 2017]
<b>Usage</b>	Research use only

