

PRODUCT INFORMATION

C-Flag&Strep Tag Tag

ACHB4 **Target Synonyms** N/A

Human ACHB4-Strep full length protein-synthetic **Description**

nanodisc **Delivery** 6~8weeks **Uniprot ID** P30926 **Expression Host HEK293**

Protein Families Ion Channels: Cys-loop Receptors

Protein Pathways N/A

Storage & Shipping

Background

The human full length ACHB4-Strep protein has a **Molecular Weight**

MW of 56.4 kDa

mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution for specific instructions. Do not use solvents with

a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

Lyophilized from nanodisc solubilization buffer (20

intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

témperature.

This gene is found within a conserved gene cluster and encodes one of the beta subunits of the nicotinic acetylcholine receptor (nAChRs) superfamily which form ligand-gated ion channels with a central pore that forms a cation channel. Neuronal nAChRs are pentameric structures that can be either homomeric or heteromeric, with heteromeric structures containing both alpha and

beta subunits. Each subunit contains an

extracellular amino terminus and four

transmembrane domains. Nicotine is one of the agonists that binds to the receptor. Variants in this gene have been associated with nicotine dependence and lung cancer. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Sep

> Email: info@dimabio.com Website: www.dimabio.com

20171

Research use only Usage Conjugate Unconjugated

