

PRODUCT INFORMATION

Tag C-Flag Tag
Target ACKR2

Synonyms CCBP2, CCR10, CCR9, CMKBR9, D6, hD6

Human ACKR2 full length protein-synthetic

nanodisc

Delivery 6~8weeks

Uniprot ID 000590

Expression Host HEK293

Formulation & Reconstitution

Background

Protein Families GPCR, Transmembrane, Druggable Genome,

Protein Pathways

GPCRDB Class A Rhodopsin-

like,Chemokines,Chemokine and Receptor,

Molecular Weight

The human full length ACKR2 protein has a MW of

43.4kDa

Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with

a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

Storage & Shipping intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

This gene encodes a beta chemokine receptor, which is predicted to be a seven transmembrane protein similar to G protein-coupled receptors. Chemokines and their receptor-mediated signal transduction are critical for the recruitment of effector immune cells to the inflammation site. This gene is expressed in a range of tissues and hemopoietic cells. The expression of this receptor in lymphatic endothelial cells and overexpression in vascular tumors suggested its function in chemokine-driven recirculation of leukocytes and

in vascular tumors suggested its function in chemokine-driven recirculation of leukocytes and possible chemokine effects on the development and growth of vascular tumors. This receptor appears to bind the majority of beta-chemokine family members; however, its specific function remains unknown. This gene is mapped to chromosome 3p21.3, a region that includes a cluster of chemokine receptor genes. [provided

by RefSeq, Jul 2008]

Usage Research use only

Conjugate Unconjugated

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com

