

PRODUCT INFORMATION

Tag	C-Flag&Strep Tag
Target	ADA2A
Synonyms	ADRA2, ADRA2R, ADRAR, ALPHA2AAR, ZNF32
Description	Human ADA2A-Strep full length protein-synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	P08913
Expression Host	HEK293
Protein Families	GPCR,Transmembrane,Druggable Genome,
Protein Pathways	GPCRDB Class A Rhodopsin-like,Monoamine GPCRs,Metabolic and Obesity,
Molecular Weight	The human full length ADA2A-Strep protein has a MW of 50.6 kDa Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
Formulation & Reconstitution	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Storage & Shipping	
Background	Alpha-2-adrenergic receptors are members of the G protein-coupled receptor superfamily. The alpha-2-adrenergic receptors are a type of adrenergic receptors (for adrenaline or epinephrine), which inhibit adenylate cyclase. These receptors include 3 highly homologous subtypes: alpha2A, alpha2B, and alpha2C. They are involved in regulating the release of neurotransmitter molecules from sympathetic nerves and from adrenergic neurons in the central nervous system. The sympathetic nervous system regulates cardiovascular function by activating adrenergic receptors in the heart, blood vessels and kidney. Studies in mouse revealed that both the alpha2A and alpha2C receptor subtypes were required for presynaptic transmitter release from the sympathetic nervous system in the heart and from central noradrenergic neurons. The alpha-2-adrenergic receptors are also involved in catecholamine signaling by extracellular regulated protein kinase 1 and 2 (ERK1/2) pathways. A clear association between the alpha-2-adrenergic receptor and disease has not been yet established. [provided by RefSeq, Sep 2019]
Usage	Research use only
Conjugate	Unconjugated

