Cat. No. FLP100132



PRODUCT INFORMATION

C-Flag Tag Tag **APLNR Target**

Synonyms AGTRL1; APJ; APJR; HG11

Human APLNR full length protein-synthetic **Description**

nanodisc **Delivery** In Stock **Uniprot ID** P35414 **Expression Host HEK293**

Storage & Shipping

Protein Families Druggable Genome, GPCR, Transmembrane

Protein Pathways Neuroactive ligand-receptor interaction

The human full length APLNR protein has a MW of **Molecular Weight**

42.7 kDa Lyophilized from nanodisc solubilization buffer (20

mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution for specific instructions. Do not use solvents with

a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

A member of the G protein-coupled receptor gene family. The encoded protein is related to the angiotensia receptor, but is actually an apelin receptor that inhibits adenylate cyclase activity and plays a counter-regulatory role against the pressure action of angiotensin II by exerting hypertensive effect. It functions in the cardiovascular and central nervous systems, in

Background

glucose metabolism, in embryonic and tumor angiogenesis and as a human immunodeficiency virus (HIV-1) coreceptor. Two transcript variants resulting from alternative splicing have been

> Email: info@dimabio.com Website: www.dimabio.com

identified.

Research use only **Usage** Conjugate Unconjugated





ELISA assay to evaluate APLNR-Nanodisc 0.2µg Human APLNR-Nanodisc per well




Figure 1. Elisa plates were pre-coated with Flag Tag APLNR-Nanodisc ($0.2\mu g/per$ well). Serial diluted anti-Flag monoclonal antibody solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Flag monoclonal antibody binding with APLNR-Nanodisc is 2.278ng/ml.

Email: info@dimabio.com Website: www.dimabio.com



Figure 2. Human APLNR-Nanodisc, Flag Tag on SDS-PAGE

