

## **PRODUCT INFORMATION**

Тад	C-Flag&Strep Tag
Target	C5AR1
Synonyms	C5A; C5AR; C5R1; CD88
Description	Human C5AR1-Strep full length protein-synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	P21730
<b>Expression Host</b>	HEK293
<b>Protein Families</b>	Druggable Genome, GPCR, Transmembrane
Protein Pathways	Complement and coagulation cascades, Neuroactive ligand-receptor interaction
Molecular Weight	The human full length C5AR1-Strep protein has a MW of 39.3 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C(Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Receptor for the chemotactic and inflammatory peptide anaphylatoxin C5a. The ligand interacts with at least two sites on the receptor: a high- affinity site on the extracellular N-terminus, and a second site in the transmembrane region which activates downstream signaling events. Receptor activation stimulates chemotaxis, granule enzyme release, intracellular calcium release and superoxide anion production
Usage	Research use only
Conjugate	Unconjugated

