

PRODUCT INFORMATION

| | |
|---|---|
| Target | CA12 |
| Synonyms | CA-XII |
| Description | Recombinant human CA12 protein with C-terminal human Fc tag |
| Delivery | In Stock |
| Uniprot ID | O43570 |
| Expression Host | HEK293 |
| Tag | C-Human Fc Tag |
| Molecular Characterization | CA12(Ala25-Ser301) hFc(Glu99-Ala330) |
| Molecular Weight | The protein has a predicted molecular mass of 57.3 kDa after removal of the signal peptide. The apparent molecular mass of CA12-hFc is approximately 55-70 kDa due to glycosylation. |
| Purity | The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining. |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | Carbonic anhydrases (CAs) are a large family of zinc metalloenzymes that catalyze the reversible hydration of carbon dioxide. They participate in a variety of biological processes, including respiration, calcification, acid-base balance, bone resorption, and the formation of aqueous humor, cerebrospinal fluid, saliva, and gastric acid. This gene product is a type I membrane protein that is highly expressed in normal tissues, such as kidney, colon and pancreas, and has been found to be overexpressed in 10% of clear cell renal carcinomas. Three transcript variants encoding different isoforms have been identified for this gene. [provided by RefSeq, Jun 2014] |
| Usage | Research use only |
| Conjugate | Unconjugated |






Figure 1. Human CA12 Protein, hFc Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

