

**PRODUCT INFORMATION**

<b>Tag</b>	C-Flag Tag
<b>Target</b>	CCG2
<b>Synonyms</b>	MRD10
<b>Description</b>	Human CCG2 full length protein-synthetic nanodisc
<b>Delivery</b>	6~8weeks
<b>Uniprot ID</b>	Q9Y698
<b>Expression Host</b>	HEK293
<b>Protein Families</b>	Ion Channels: Other
<b>Protein Pathways</b>	N/A
<b>Molecular Weight</b>	The human full length CCG2 protein has a MW of 36kDa
<b>Formulation &amp; Reconstitution</b>	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	The protein encoded by this gene is a type I transmembrane AMPA receptor regulatory protein (TARP). TARPs regulate both trafficking and channel gating of the AMPA receptors. The AMPA subtype of ionotropic glutamate receptors are ligand gated ion channels that are typically activated by glutamate released from presynaptic neuron terminals and mediate fast neurotransmission in excitatory synapses. TARPs thus play an important role in synaptic plasticity, learning and memory. Mutations in this gene cause an autosomal dominant form of cognitive disability. [provided by RefSeq, Jul 2017]
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated

