

PRODUCT INFORMATION

Target	CCR6
Synonyms	BN-1;C-C CKR-6;CC-CKR-6;CCR-6;CD196;CKR-L3;CKRL3;CMKBR6;DCR2;DRY6;GPR29;GPRCY4;STRL22
Description	Recombinant Human CCR6 with C-terminal human Fc tag
Delivery	In Stock
Uniprot ID	P51684
Expression Host	HEK293
Tag	C-Human Fc Tag
Molecular Characterization	CCR6(Met1-Leu47) hFc(Glu99-Ala330)
Molecular Weight	The protein has a predicted molecular mass of 31.6 kDa after removal of the signal peptide. The apparent molecular mass of CCR6-hFc is approximately 35-55 kDa due to glycosylation.
Purity	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene encodes a member of the beta chemokine receptor family, which is predicted to be a seven transmembrane protein similar to G protein-coupled receptors. The gene is preferentially expressed by immature dendritic cells and memory T cells. The ligand of this receptor is macrophage inflammatory protein 3 alpha (MIP-3 alpha). This receptor has been shown to be important for B-lineage maturation and antigen-driven B-cell differentiation, and it may regulate the migration and recruitment of dendritic and T cells during inflammatory and immunological responses. Alternatively spliced transcript variants that encode the same protein have been described for this gene. [provided by RefSeq, Jul 2008]
Usage	Research use only
Conjugate	Unconjugated




Figure 1. Human CCR6 Protein, hFc Tag on SDS-PAGE under reducing condition.
 Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China
 Telephone: +1 2409940618(USA)
 /+86-18062749453(China)
 /+86-400-006-0995(China)

Email: info@dimabio.com
 Website: www.dimabio.com



DIMABIO CONFIDENTIAL

