

**PRODUCT INFORMATION**

<b>Target</b>	CD112
<b>Synonyms</b>	NECTIN2;HVEB;PRR2;PVRL2;PVRR2
<b>Description</b>	Recombinant Human CD112 with C-terminal 6×His tag
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	Q92692
<b>Expression Host</b>	HEK293
<b>Tag</b>	C-6×His Tag
<b>Molecular Characterization</b>	CD112(Gln32-Gly360) 6×His tag
<b>Molecular Weight</b>	The protein has a predicted molecular mass of 36.1 kDa after removal of the signal peptide. The apparent molecular mass of CD112-His is approximately 35-55 kDa due to glycosylation.
<b>Purity</b>	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	This gene encodes a single-pass type I membrane glycoprotein with two Ig-like C2-type domains and an Ig-like V-type domain. This protein is one of the plasma membrane components of adherens junctions. It also serves as an entry for certain mutant strains of herpes simplex virus and pseudorabies virus, and it is involved in cell to cell spreading of these viruses. Variations in this gene have been associated with differences in the severity of multiple sclerosis. Alternate transcriptional splice variants, encoding different isoforms, have been characterized.
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated






Figure 1. Human CD112 Protein, His Tag on SDS-PAGE under reducing condition.

### Human CD112,His Tagged protein ELISA

0.2  $\mu$ g of Human CD112, His tagged protein per well




Figure 2. ELISA plate pre-coated by 2  $\mu$ g/mL (100  $\mu$ L/well) Human CD112 Protein, His Tag(PME100576) can bind Anti-CD112 antibody(DMC224), IgG1 Chimeric mAb in a linear range of 0.64-16.00 ng/mL.

