

PRODUCT INFORMATION

| | |
|---|---|
| Target | CD32a |
| Synonyms | CD32;CD32A;CDw32;FCG2;FcGR;FCGR2;FCGR2A1;IGFR2 |
| Description | Recombinant Human CD32A(167H) with C-terminal human Fc tag |
| Delivery | In Stock |
| Uniprot ID | P12318 |
| Expression Host | HEK293 |
| Tag | C-Human Fc Tag |
| Molecular Characterization | CD32a(H167)(Ala36-Ile218) hFc(Glu99-Ala330) |
| Molecular Weight | The protein has a predicted molecular mass of 46.4 kDa after removal of the signal peptide. The apparent molecular mass of CD32A(167H)-hFc is approximately 55-70 kDa due to glycosylation. |
| Purity | The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining. |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Storage & Shipping | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | This gene encodes one member of a family of immunoglobulin Fc receptor genes found on the surface of many immune response cells. The protein encoded by this gene is a cell surface receptor found on phagocytic cells such as macrophages and neutrophils, and is involved in the process of phagocytosis and clearing of immune complexes. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Oct 2008] |
| Usage | Research use only |
| Conjugate | Unconjugated |






Figure 1. Human CD32a (H167) Protein, hFc Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

