

PRODUCT INFORMATION

CD45 Target

B220;CD45;CD45R;GP180;L-CA;LCA;LY5;T200 **Synonyms**

Recombinant Human CD45 with C-terminal 6×His **Description**

In Stock **Delivery Uniprot ID** P08575 **Expression Host HEK293** C-6×His Tag Tag

Molecular

Purity

Reconstitution

Background

CD45(Gln26-Lys577) 6×His tag Characterization

The protein has a predicted molecular mass of

61.6 kDa after removal of the signal peptide. The apparent molecular mass of CD45-His is **Molecular Weight**

approximately 35-55 kDa due to glycosylation.

The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % Formulation &

- 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that

regulate a variety of cellular processes including cell growth, differentiation, mitosis, and oncogenic transformation. This PTP contains an extracellular domain, a single transmembrane segment and two tandem intracytoplasmic

catalytic domains, and thus is classified as a receptor type PTP. This PTP has been shown to be an essential regulator of T- and B-cell antigen receptor signaling. It functions through either direct interaction with components of the antigen receptor complexes, or by activating various Src family kinases required for the antigen receptor signaling. This PTP also suppresses JAK kinases, and thus pinceling as a regulator of cytokine

receptor signaling. Alternatively spliced transcripts variants of this gene, which encode distinct isoforms, have been reported. [provided

> Email: info@dimabio.com Website: www.dimabio.com

by RefSeq, Jun 2012]

Usage Research use only

Conjugate Unconjugated

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)








Figure 1. Human CD45 Protein, His Tag on SDS-PAGE under reducing condition.

Human CD45, His Tagged protein ELISA

0.2 μg of Human CD45, HIs tagged protein per well




Figure 2. ELISA plate pre-coated by 2 μ g/mL (100 μ L/well) Human CD45 Protein, His Tag(PME100581) can bind Anti-CD45 antibody(DM193), Rabbit mAb in a linear range of 0.64–16.00 ng/mL.

