

## **PRODUCT INFORMATION**

CD62E **Target** 

**Synonyms** ELAM; ESEL; SELE; ELAM1; LECAM2; selectin-e Recombinant human CD62E Protein with C-**Description** 

terminal 6×His tag

**Delivery** In Stock **Uniprot ID** P16581 **Expression Host HEK293** Tag C-6×His Tag

Molecular

Storage & Shipping

**Background** 

Purity

CD62E(Trp22-Pro556) 6×His tag Characterization

The protein has a predicted molecular mass of **Molecular Weight** 

59.5 kDa after removal of the signal peptide. The apparent molecular mass of CD62E-His is

approximately 100-130 kDa due to glycosylation.

The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue

staining.

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

The protein encoded by this gene is found in cytokine-stimulated endothelial cells and is thought to be responsible for the accumulation of blood leukocytes at sites of inflammation by mediating the adhesion of cells to the vascular lining. It exhibits structural features such as the presence of lectin- and EGF-like domains followed by short consensus repeat (SCR) domains that contain 6 conserved cysteine residues. These

proteins are part of the selectin family of cell adhesion molecules. Adhesion molecules participate in the interaction between leukocytes and the endothelium and appear to be involved in the pathogenesis of atherosclerosis. [provided by

RefSeq, Jul 2008]

Usage Research use only Conjugate Unconjugated









Figure 1. Human CD62E Protein, His Tag on SDS-PAGE under reducing condition.



