

PRODUCT INFORMATION

Target	CEACAM1
Synonyms	BGP;BGP1;BGPI
Description	Recombinant human CEACAM1 protein with C-terminal 6×His tag
Delivery	In Stock
Uniprot ID	P13688
Expression Host	HEK293
Tag	C-6×His Tag
Molecular Characterization	CEACAM1(Gln35-Gly428) 6×His tag
Molecular Weight	The protein has a predicted molecular mass of 44.1 kDa after removal of the signal peptide. The apparent molecular mass of CEACAM1-His is approximately 70-100 kDa due to glycosylation.
Purity	The purity of the protein is greater than 85% as determined by SDS-PAGE and Coomassie blue staining.
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene encodes a member of the carcinoembryonic antigen (CEA) gene family, which belongs to the immunoglobulin superfamily. Two subgroups of the CEA family, the CEA cell adhesion molecules and the pregnancy-specific glycoproteins, are located within a 1.2 Mb cluster on the long arm of chromosome 19. Eleven pseudogenes of the CEA cell adhesion molecule subgroup are also found in the cluster. The encoded protein was originally described in bile ducts of liver as biliary glycoprotein. Subsequently, it was found to be a cell-cell adhesion molecule detected on leukocytes, epithelia, and endothelia. The encoded protein mediates cell adhesion via homophilic as well as heterophilic binding to other proteins of the subgroup. Multiple cellular activities have been attributed to the encoded protein, including roles in the differentiation and arrangement of tissue three-dimensional structure, angiogenesis, apoptosis, tumor suppression, metastasis, and the modulation of innate and adaptive immune responses. Multiple transcript variants encoding different isoforms have been reported, but the full-length nature of all variants has not been defined. [provided by RefSeq, May 2010]
Usage	Research use only
Conjugate	Unconjugated






Figure 1. Human CEACAM1 Protein, His Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

