

PRODUCT INFORMATION

Tag	C-Flag&Strep Tag
Target	CELR1
Synonyms	ADGRC1, CDHF9, FMI2, HFMI2, LMPHM9, ME2
Description	Human CELR1-Strep full length protein-synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	Q9NYQ6
Expression Host	HEK293
Protein Families	Transmembrane,Druggable Genome,
Protein Pathways	GPCRDB Other,
Molecular Weight	The human full length CELR1-Strep protein has a MW of 329.5 kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	The protein encoded by this gene is a member of the flamingo subfamily, part of the cadherin superfamily. The flamingo subfamily consists of nonclassic-type cadherins; a subpopulation that does not interact with catenins. The flamingo cadherins are located at the plasma membrane and have nine cadherin domains, seven epidermal growth factor-like repeats and two laminin A G-type repeats in their ectodomain. They also have seven transmembrane domains, a characteristic unique to this subfamily. It is postulated that these proteins are receptors involved in contact-mediated communication, with cadherin domains acting as homophilic binding regions and the EGF-like domains involved in cell adhesion and receptor-ligand interactions. This particular member is a developmentally regulated, neural-specific gene which plays an unspecified role in early embryogenesis. [provided by RefSeq, Jul 2008]
Usage	Research use only
Conjugate	Unconjugated

