

## PRODUCT INFORMATION

<b>Target</b>	CGRPR
<b>Synonyms</b>	CRLR; CALCRL; LMPHM8
<b>Description</b>	Recombinant human CGRPR Protein with C-terminal human Fc tag
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	Q16602
<b>Expression Host</b>	HEK293
<b>Tag</b>	C-Human Fc tag
<b>Molecular Characterization</b>	CGRPR(Glu23-Asn140) hFc(Glu99-Ala330)
<b>Molecular Weight</b>	The protein has a predicted molecular mass of 39.9 kDa after removal of the signal peptide. The apparent molecular mass of CGRPR-hFc is approximately 35-70 kDa due to glycosylation.
<b>Purity</b>	The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	Enables adrenomedullin binding activity; adrenomedullin receptor activity; and calcitonin gene-related peptide receptor activity. Involved in several processes, including G protein-coupled receptor signaling pathway; cellular response to sucrose stimulus; and receptor internalization. Located in endoplasmic reticulum; endosome; and lysosome. Part of CGRP receptor complex and adrenomedullin receptor complex. Colocalizes with plasma membrane. Implicated in hereditary lymphedema. [provided by Alliance of Genome Resources, Apr 2022]
<b>Usage</b>	Research use only





Figure 1. Human CGRPR Protein, hFc Tag on SDS-PAGE under reducing condition.

DIMABIO CONFIDENTIAL

