

PRODUCT INFORMATION

Tag	C-Flag Tag
Target	CLCA2
Synonyms	CACC, CACC3, CLCRG2, CaCC-3
Description	Human CLCA2 full length protein-synthetic nanodisc
Delivery	6~8weeks
Uniprot ID	Q9UQC9
Expression Host	HEK293
Protein Families	Ion Channels: Other
Protein Pathways	N/A
Molecular Weight	The human full length CLCA2 protein has a MW of 103.9kDa
Formulation & Reconstitution	Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Do not use solvents with a pH below 6.5 or those containing high concentrations of divalent metal ions (greater than 5 mM) in subsequent experiments.
Storage & Shipping	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	This gene encodes a member of the calcium-activated chloride channel regulator (CLCR) family of proteins. Members of this family regulate the transport of chloride across the plasma membrane. The encoded protein is autoproteolytically processed to generate N- and C- terminal fragments. Expression of this gene is upregulated by the tumor suppressor protein p53 in response to DNA damage. In breast cancer, expression of this gene is downregulated and the encoded protein may inhibit migration and invasion while promoting mesenchymal-to-epithelial transition in cancer cell lines. [provided by RefSeq, Sep 2016]
Usage	Research use only
Conjugate	Unconjugated

